

Compensating for a Bite Block in /t/ and /d/ in French : Electropalatography Study

Sandrine Clairet

Université de Provence

E-mail: sandrine.clairet@univ-provence.fr

Abstract

This paper deals with various movements of the tongue in perturbed speech. The activity of the jaw has been neutralized by mean of bite block in order to observe immediate compensatory tongue movement when the jaw was blocked. Electropalatography was used to monitor linguopalatal contact patterns in French /t/ and /d/ in normal and in bite-block speech. We thus observed how spatial and temporal changes of the tongue configurations palliate the immobilization of the mandible. Results showed that the presence of bite block immediatly modified the duration and the amplitude of the complete closure and effects of bite block were speaker dependent. Linguo-palatal contacts were made further forward than in normal speech, and reached a different articulatory target.

1 Introduction

Studies on oral-articulatory perturbations have provided valuable insights into motor equivalence, or how different muscle activation patterns can achieve the same articulatory or acoustic goals. It's well known that functional dependences exist between the jaw and the tongue which together formed part of a coordinative structure [1] [2] [3]. So articulatory compensations could be immediatly achieved as a consequence of this inter-articulatory coordination [1]. In particular, previous studies on vowels of steady-state structural perturbations caused by the insertion of a bite block into the mouth have revealed that articulatory compensation made by adjusting articulator positions allowed the phoneme-specific acoustic goal to be achieved [4] [5] [6].

But, few works have studied the production of consonant with bite blocks even if there is somewhat less flexibility in consonant production as compared to the production of vowels. Moreover certain consonants require greater articulatory precision than others. Some authors have studied the production of /t/ and /s/ in English and in Arabic languages, in normal and bite block speech and their articulatory and perceptual results suggested the speakers did not compensated completely. Compensation was not instantaneous and some information pertaining to specific parameter values

were encoded in central representations [6] [7]. The original approach of the present study consists of producing French consonants /t/ and /d/ with bite block and comparing our results with those made in the same condition, but in other languages. We used the paradigm of bite block speech to infer compensatory movements of the tongue, thus reaching other articulatory target.

Previous results of an articulatory study showed that bite block induced different spatio-temporal modifications: although a necessary constriction was achieved, we could note a reduction as well as a reinforcement of the general movement of the tongue. See [8] for details.

Therefore, the aim of this paper is to study the spatio temporal modifications of the medial phase of the consonant, specifically on the complete closure (defined as the total obstruction of the oral airflow) and its location which can be free to vary within the antero-posterior dimension : the bite block are not only expected to reduce the area of tongue palate contact but to shift forward the place of articulation.

2 Method

The corpus was composed of Consonant-Vowel-Consonant (CVC) sequences embedded in the carrier phrase: "Il ne dit pas CVC encore" (He doesn't say CVC again). The items studied were C1V syllables containing /t/ and /d/ stops, alternately /a/ and /i/ respectively the most open and closed vowels in the French vocalic system, repeated twelve times by two French speakers, in normal and bite block speech.

To fix the jaw, we fashioned acrylic bite blocks with a thermo malleable dentistry paste, making individual indentations for tooth cups to prevent slipping. When held between the molar teeth, bite blocks were adjusted to ensure a vertical distance of 17 mm measured between the central incisors for both speakers. Bite block size was chosen based on previous researches with bite blocks.

Recordings were made using the electropalatography (EPG) system of Reading [10] which allowed a spatio temporal analysis of the linguo palatal (LP) contacts. Even though both speakers were used to wearing artificial palate and making recordings with, we allowed a thirty minute

adaptation period during the calibration of the recording system. However, recordings were made immediately after the insertion of the bite block into the mouth. EPG used an acrylic 1,5 mm thick palate created from a molding of the hard palate of our speakers. Sixty two contact electrodes were placed in eight rows, six on the first anterior row and eight on the others. The articulation zone of alveolar stop consonants was empirically defined as being the most anterior area of the palate represented by the four first anterior rows of the hard palate [10]. Measures were collected and based on a manual segmentation of the temporal evolution curve of the LP contacts coupled with the LP contact patterns of each EPG frames, the both simultaneously with the acoustic signal.

Articulatory parameters were taken on the complete closure. Complete closure duration was defined as the interval between first total closure (the first EPG frame showing a complete row contacted) and closure release (the first EPG frame showing an opening in the complete row contacted). It's important to note that in bite block speech, the mean rate of measurable complete closure was the good 88% score for both speakers. Of course, in this case, the main obstacle to successful measurement was lack of the total closure on the EPG patterns. The maximum number of LP contacts was taken during this interval. In order to detect the distribution of the LP contacts, the center of gravity was calculated on the antero-posterior axis of the palate. The axis of reference was the median horizontal limit between the four anterior rows (-1 à -4) and the four posterior rows (+1 à +4). So the further the center of gravity receded, the more we could observe a negative value. This factor was relevant to inform us on changes in the area of articulation.

3 Results

3.1 Duration of the complete closure




Figure 1: Mean duration relative to the entire consonant

As can be seen in Figure 1 which shows the mean duration of the closure relative to the entire consonant, the effect of the bite block was speaker dependant. For speaker B, closure duration had a tendency to be significantly shorter in perturbed speech than in normal speech and the influence of the bite block was consequent [F (3,161)=4,492; p=0,0047]. The mean difference between the bite block and normal speech samples was -2,96 %. For speaker Y, the duration of closure was lengthened of 4,70% in bite block speech, excepted for the sequence /dad/ which presented a decrease of -2,59 %. However, for this speaker, it seems that the effects of the bite block had no significant influence on the duration of the closure [F (3,167)=2,355; p=0,073].

3.2 Amplitude: Number of LP contacts




Figure 2: Amplitude of LP contacts

Speaker B contacted fewer sensors in bite block speech than in normal speech in all sequences. There was a mean difference of -6,52 sensors contacted between perturbed and normal speech. The /dad/ sequence was the most affected by the bite block as it lost -10,66. For this speaker, amplitude of contacts was reduced with block. Indeed, there was a direct relationship between the influence of the bite block and the number of LP contacts [F (3,175)=53,44; p<0,0001]. Speaker Y tended to contact more sensors in the bite block condition than in the normal condition, apart from the /dad/ sequence. The lowest mean difference between perturbed and normal speech was of +2,42 sensors contacted.

Although the bite block made have varied the duration and the amplitude, variations differed strongly according to the speakers ; not including the /dad/ sequence for which we noted a similar spatial reduction of the amplitude and shortening of the complete closure for the both speakers.

speakers contacted more sensors in bite block speech than in normal speech.[6]. Certainly, a part of these differences between subjects can be explained by dissimilarities in implementing consonant in their native language. But another part of these differences is due to the existence of individual strategies for compensation depending on a set of personal factors such as morphology, articulatory capacities, experiences, knowledge...

These results raise the questions of whether the articulatory variations resulting from the use of the bite block caused significant acoustic differences. Currently, we are making a study of the acoustic parameters of /t/ and /d/ in normal and bite block speech. Spectral characteristics of the burst noise are measured in order to compare them to the articulatory parameters. The acoustic study should confirm that an acceptable acoustic target may be reached in different articulatory manners. Shortly we will be able to add argument to demonstrate that the variability in speech can be explained by the existence of motor equivalences that control the articulatory movements in speech [11] and this control must be adapted 'on line' with a large set of constraints of production.

5 References

- [1] J. Kelso & al. Functionally-specific cooperation following jaw perturbation during speech: Evidence for coordinative structures, *Journal of Experimental Psychology: Human Perception and Performance*, 10, pp. 812-832, 1984.
- [2] J.S., Perkell & al. Trading Relations Between Tongue Body Raising and Lip Rounding in Production of the Vowel /u/: a Pilot "Motor Equivalence" Study, *Journal of the Acoustical Society of America*, 93(5), 2948-2961, 1993.
- [3] P. Keating & al. Variability in jaw height for segments in English and Swedish VCV, *Journal of Phonetics*, 22, pp407-422, 1994.
- [4] B.Lindblom & al. Formant frequencies of some fixed mandible vowels and a model of speech motor programming by predictive simulation, *Journal of Phonetics*, 7, pp.147-161, 1979.
- [5] C.A. Fowler & M. Turvey Immediate compensation in bite block speech, *Phonetica* 37, pp. 306-326, 1980.
- [6] J. E.. Flege & al. Compensating for a bite block in /s/ and /t/ production: palatographic, acoustic et perceptual data, *Journal of the Acoustical Society of America*, 83(1), pp. 212-228, 1988.
- [7] D. McFarland & S. Baum Incomplète compensation to articulatory perturbation, *Journal of the Acoustical Society of America*, 97(3), pp. 1865-1873, 1995.
- [8] Clairret, S. Compensation articuloire dans la production des occlusives linguales du français, in *Perturbations et Réajustements : langue et langage*, Publicationse l'Université de Strasbourg, B.
- Vaxelaire, R.Sock, G. Kleiber, F. Marsac (Eds.), pp.51-62, 2007.
- [9] W.J. Hardcastle & al. EPG Data Reduction Methods and Their Implications for Studies of Lingual Coarticulation, *Journal of Phonetic*, 19, 251-266, 1991.
- [10] P. Ladefoged & I.Maddieson, the sounds of the world's languages, *Blackwell Publishers*, 19996
- [11] J.H. Abbs & V.L.Gracco, Control of complex motor gestures: orofacial muscle responses to load perturbation of lip during speech, *Journal of neurophysiology*, 51, pp705-723, 1984.